

## APPLIED MATHEMATICS - HIGHER LEVEL

FRIDAY, 28 JUNE - AFTERNOON, 2.00 to 4.30

Six questions to be answered. All questions carry equal marks.  
 Mathematics Tables may be obtained from the Superintendent.  
 Take the value of  $g$  to be  $9.8 \text{ metres/second}^2$ .  $\vec{i}$  and  $\vec{j}$  are perpendicular unit vectors.

1. A bus  $12.5 \text{ m}$  long travels with constant acceleration. The front of the bus passes a point,  $p$ , with speed  $u$  while the rear of the bus passes  $p$  with speed  $v$ .


Find, in terms of  $u$  and  $v$

- the time taken by the bus to pass  $p$ .
- what fraction of the length of the bus passes the point  $p$  in half this time.

2. A particle is projected from a point  $o$  with initial velocity  $u$ , up a plane inclined at an angle of  $60^\circ$  to the horizontal. The direction of projection makes an angle  $\theta$  with the inclined plane. (The plane of projection is vertical and contains the line of greatest slope). The maximum height reached above the inclined plane is  $H$ .

- Express
- the velocity and displacement from  $o$  of the particle after  $t$  seconds, in terms of  $\vec{i}$  and  $\vec{j}$ , where these are the unit vectors along and perpendicular to the plane, respectively.
  - $H$ , in terms of  $u$  and  $\theta$ .
  - the time interval, in terms of  $\sin 2\theta$ , between the instants when the particle is at a height,  $H \sin^2 \theta$ , above the inclined plane.

3. Two blocks  $A$  and  $B$  have masses  $2 \text{ kg}$  and  $x \text{ kg}$ , respectively. They are connected by a string and slide down an inclined plane which makes an angle  $\sin^{-1}(\frac{3}{5})$  with the horizontal. The coefficient of friction between  $A$  and the plane is  $\frac{1}{4}$ , and between  $B$  and the plane is  $\frac{1}{2}$ .



- Show in a diagram the forces acting on each block when the system is released from rest.
  - Find the acceleration  $f$  of the system in terms of  $x$ .
  - For what value of  $x$  would the acceleration of the blocks be  $0.9f$ ?
4. State the laws governing the oblique collision of elastic spheres.  
 A smooth sphere  $A$  impinges obliquely on an identical smooth sphere  $B$  which is at rest. The direction of  $A$  before and after impact makes angles  $60^\circ$  and  $\theta$ , respectively, with the line of centres.

- Prove that  $\tan \theta = \frac{2\sqrt{3}}{1-e}$  where  $e$  is the coefficient of restitution between the spheres.
- Show that the maximum percentage loss in kinetic energy due to the impact is  $12\frac{1}{2}\%$ .
- For what value of  $e$  will the kinetic energies of  $A$  and  $B$  after impact be in the ratio  $7 : 1$ ?


5. (a) Two cars  $A$  and  $B$  are moving along straight roads which are at right angles to each other, with uniform velocities  $3 \text{ m/s}$  and  $4 \text{ m/s}$ , respectively. When  $B$  is at the crossroads,  $A$  is  $100 \text{ m}$  away. Calculate the time interval for which the distance between the cars is not greater than  $82 \text{ m}$ .
- (b) A car of mass  $750 \text{ kg}$  attains a maximum speed of  $30 \text{ m/s}$  when travelling down an incline of  $1$  in  $25$  with the engine switched off. It can attain a maximum speed of  $20 \text{ m/s}$  up the same incline when the engine is working. The resistance to motion in each case is proportional to the square of the speed. Find
- the power at which the engine is working
  - the maximum speed of the car along a level road, if it works at the same power and its resistance is again proportional to the square of the speed.

6. Define simple harmonic motion.

A body of mass  $0.25 \text{ kg}$  hangs from a spiral spring. When pulled down  $10 \text{ cm}$  below its equilibrium position and released, it vibrates with simple harmonic motion of period  $2 \text{ s}$ .

- Find its velocity as it passes through the equilibrium position.
- What is the shortest time taken to travel from a point  $2 \text{ cm}$  below the equilibrium position to a point  $2 \text{ cm}$  above the equilibrium position?
- Find the elastic constant of the spring.
- By how much will the spring shorten when the body is removed?


7. One end of a uniform metre stick of mass  $m$  is placed against a vertical wall. The other end is held by a light inelastic string making an angle  $\theta$  with the metre stick. The coefficient of friction,  $\mu$ , between the end of the metre stick and the wall is  $0.4$ .



- Show in a diagram the forces acting on the metre stick.
- Show that if the metre stick is to remain in equilibrium the maximum value of  $\theta$  is given by  $\tan \theta = \mu$ .
- If a mass  $m$  is suspended from the metre stick at a distance  $x$  from the wall, show that the stick is on the point of slipping when


$$\tan \theta = \frac{2(1 + 2x)}{5(3 - 2x)}$$

8. A uniform square lamina  $abcd$ , of mass  $3m$  and side  $\sqrt{2}$ , is free to rotate with its plane vertical about a smooth horizontal axis through a point  $p$  on the line  $ac$ . A mass  $m$  is attached at each of the points  $a$  and  $c$ .



- (i) If  $|ap| = 1 - x$ , prove that the moment of inertia of the system about a horizontal axis through  $p$  is  $m(3 + 5x^2)$ .
- (ii) If the system oscillates about  $p$ , find in terms of  $x$ , the period for small oscillations.
- (iii) Find the value of  $x$  which gives the minimum period when oscillations are small.
9. (a) A piece of gold-aluminium alloy of mass  $10$  kg, weighs  $72$  N in water. If the relative densities of gold and aluminium are  $19.6$  and  $2.45$  respectively, find
- the relative density of the alloy
  - the mass of each metal in the alloy
  - what fraction of the total volume of the alloy is gold.

- (b) A uniform rod of relative density  $0.25$  is free to turn about its lower end, which is fixed at a depth  $0.4$  m in water. The rod is in equilibrium when partially immersed and making an angle of  $60^\circ$  with the vertical. Find the length of the rod.



10. (a) Find the solution of the differential equation

$$3y^2(x - 1) \frac{dy}{dx} = 1 + y^3$$

if  $y = 0$  when  $x = 2$ .

- (b) A particle of mass  $m$  moves in a straight line. The only force acting on it being a resistance  $mkv^2$ , where  $v$  is its speed and  $k$  is a constant. It is initially projected from the point  $o$  with speed  $u$ . When the particle reaches a point  $p$  on the line its speed is  $u/3$ .
- Show that the average speed between  $o$  and  $p$  is  $\frac{1}{2}u \log 3$ .
  - Find the speed of the particle when it is at the midpoint of  $[op]$ .